



# Mark Scheme (Results)

Summer 2024

Pearson Edexcel International Advanced Level  
In Pure Mathematics (WMA11) Paper 01

## **Edexcel and BTEC Qualifications**

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at [www.edexcel.com](http://www.edexcel.com) or [www.btec.co.uk](http://www.btec.co.uk). Alternatively, you can get in touch with us using the details on our contact us page at [www.edexcel.com/contactus](http://www.edexcel.com/contactus).

## **Pearson: helping people progress, everywhere**

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: [www.pearson.com/uk](http://www.pearson.com/uk)

Summer 2024

Question Paper Log Number P75707A

Publications Code WMA11\_01\_2406\_MS

All the material in this publication is copyright

© Pearson Education Ltd 2024

## General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

## General Instructions for Marking

1. The total number of marks for the paper is 75.
2. The Edexcel Mathematics mark schemes use the following types of marks:
  - **M** marks: Method marks are awarded for 'knowing a method and attempting to apply it', unless otherwise indicated.
  - **A** marks: Accuracy marks can only be awarded if the relevant method (M) marks have been earned.
  - **B** marks are unconditional accuracy marks (independent of M marks)

Marks should not be subdivided.

### 3. Abbreviations

These are some of the traditional marking abbreviations that will appear in the mark schemes and can be used if you are using the annotation facility on ePEN:

- bod – benefit of doubt
- ft – follow through
  - the symbol  $\checkmark$  will be used for correct ft
- cao – correct answer only
- cso – correct solution only. There must be no errors in this part of the question to obtain this mark
- isw – ignore subsequent working
- awrt – answers which round to
- SC – special case
- oe – or equivalent (and appropriate)
- d... or dep – dependent
- indep – independent
- dp – decimal places
- sf – significant figures
- \* – The answer is printed on the paper or ag- answer given
- $\square$  or d... – The second mark is dependent on gaining the first mark

4. All A marks are 'correct answer only' (cao), unless shown, for example, as A1 ft to indicate that previous wrong working is to be followed through. After a misread however, the subsequent A marks affected are treated as A ft, but manifestly absurd answers should never be awarded A marks.

5. For misreading which does not alter the character of a question or materially simplify it, deduct two from any A or B marks gained, in that part of the question affected. If you are using the annotation facility on ePEN, indicate this action by 'MR' in the body of the script.
6. If a candidate makes more than one attempt at any question:
  - a) If all but one attempt is crossed out, mark the attempt which is NOT crossed out.
  - b) If either all attempts are crossed out or none are crossed out, mark all the attempts and score the highest single attempt.
7. Ignore wrong working or incorrect statements following a correct answer.

## General Principles for Pure Mathematics Marking

(NB specific mark schemes may sometimes override these general principles)

### Method mark for solving 3 term quadratic:

#### 1. Factorisation

$(x^2 + bx + c) = (x + p)(x + q)$ , where  $|pq| = |c|$  leading to  $x = \dots$

$(ax^2 + bx + c) = (mx + p)(nx + q)$ , where  $|pq| = |c|$  and  $|mn| = |a|$  leading to  $x = \dots$

#### 2. Formula

Attempt to use correct formula (with values for a, b and c)

#### 3. Completing the square

Solving  $x^2 + bx + c = 0$ :  $(x \pm \frac{b}{2})^2 \pm q \pm c$ ,  $q \neq 0$  leading to  $x = \dots$

### Method marks for differentiation and integration:

#### 1. Differentiation

Power of at least one term decreased by 1 ( $x^n \rightarrow x^{n-1}$ )

#### 2. Integration

Power of at least one term increased by 1 ( $x^n \rightarrow x^{n+1}$ )

### Use of a formula

Where a method involves using a formula that has been learnt, the advice given in recent examiners' reports is that the formula should be quoted first.

### Normal marking procedure is as follows:

Method mark for quoting a correct formula and attempting to use it, even if there are small mistakes in the substitution of values.

Where the formula is not quoted, the method mark can be gained by implication from correct working with values, but may be lost if there is any mistake in the working.

### Exact answers

Examiners' reports have emphasised that where, for example, an exact answer is asked for, or working with surds is clearly required, marks will normally be lost if the candidate resorts to using rounded decimals.

### Answers without working

The rubric says that these may not gain full credit. Individual mark schemes will give details of what happens in particular cases. General policy is that if it could be done "in your head", detailed working would not be required. Most candidates do show working, but there are occasional awkward cases and if the mark scheme does not cover this, please contact your team leader for advice.

Question	Scheme	Marks
<b>1</b>	$\int \left( 10x^4 - \frac{3}{2x^3} - 7 \right) dx = 2x^5 + \frac{3}{4}x^{-2} - 7x + c$	M1A1A1
		<b>(3)</b>
		<b>Total 3</b>

**Notes:**

**M1:** For increasing one of the powers of  $x$  by 1 e.g.  $x^4 \rightarrow \dots x^5$  or  $x^{-3} \rightarrow \dots x^{-2}$  or  $7 \rightarrow \dots 7x$

**A1:** Any **two** correctly integrated terms, unsimplified or simplified.

Allow e.g.  $\frac{10}{4+1}x^{4+1}$ ,  $-\frac{3}{-4}x^{-3+1}$ ,  $-7x^1$

**A1:**  $2x^5 + \frac{3}{4}x^{-2} - 7x + c$  all correct and simplified in one expression and including “+  $c$ ”.

Allow correct simplified equivalents for  $\frac{3}{4}x^{-2}$  e.g.  $0.75x^{-2}$ ,  $\frac{3}{4x^2}$  but do **not** allow  $-7x^1$  for  $-7x$  or  $\frac{2}{1}$  for 2

Award this mark once a correct simplified expression is seen and apply isw if necessary.

Condone poor notation e.g. spurious integral signs, “ $dx$ ”,  $\frac{dy}{dx} = \dots$  and look for the correct expression.

Question	Scheme	Marks
2(i)(a)	$2^{n+3} = 2^n \times 2^3 = 8m$	B1
		(1)
(b)	$16^{3n} = (2^4)^{3n}$	M1
	$= 2^{12n} = (2^n)^{12} = m^{12}$	A1
		(2)

**Notes:**

(i)(a)

**B1:** For  $8m$  (Condone 8M). Do **not** allow  $2^3m$ .

(b)

**M1:** For writing  $16^{3n}$  correctly as an expression involving a power of 2.

Examples:  $(2^4)^{3n}$ ,  $(2^2)^{6n}$ ,  $(2^{3n})^4$ ,  $(2^n)^{12}$ ,  $(2^{12})^n$ ,  $2^{12n}$

**A1:** For  $m^{12}$  (Condone  $M^{12}$ )

Correct answer only scores both marks.



<b>(ii)</b>	$x\sqrt{3}-3 = x+\sqrt{3} \Rightarrow x\sqrt{3}-x = 3+\sqrt{3} \Rightarrow x(\sqrt{3}-1) = 3+\sqrt{3} \Rightarrow x = \dots$	M1
	$= \frac{3+\sqrt{3}}{\sqrt{3}-1} \times \frac{\pm(\sqrt{3}+1)}{\pm(\sqrt{3}+1)}$	M1
	$= \frac{\pm(4\sqrt{3}+6)}{\pm(3-1)} = 3+2\sqrt{3}$	A1
		<b>(3)</b>

**Notes:**

**In this part, the A1 depends on both M marks**

(ii)

**M1:** Collects  $x$  terms to one side, factorises and makes  $x$  the subject

It is for processing the given equation and reaching  $x = \frac{\alpha + \beta\sqrt{3}}{\gamma + \delta\sqrt{3}}$ ,  $\alpha, \beta, \gamma, \delta \neq 0$

**M1:** Correct attempt to rationalise their denominator.

Not formally dependent but requires  $\frac{\dots}{\gamma + \delta\sqrt{3}} = \frac{\dots}{\gamma + \delta\sqrt{3}} \times \frac{\gamma - \delta\sqrt{3}}{\gamma - \delta\sqrt{3}}$  or equivalent.

Sight of e.g.  $\frac{3+\sqrt{3}}{\sqrt{3}-1} \times \frac{\sqrt{3}+1}{\sqrt{3}+1}$  is sufficient for this mark **but may be implied** by e.g.

$$\frac{3+\sqrt{3}}{\sqrt{3}-1} = \frac{3\sqrt{3}+3+3+\sqrt{3}}{3-1}$$

**A1:** For  $3+2\sqrt{3}$  with at least one intermediate step e.g.

$$\frac{3+\sqrt{3}}{\sqrt{3}-1} = \frac{3\sqrt{3}+3+3+\sqrt{3}}{3-1} \quad \text{or} \quad \frac{3+\sqrt{3}}{\sqrt{3}-1} \times \frac{\sqrt{3}+1}{\sqrt{3}+1} = \frac{4\sqrt{3}+6}{2} \quad \text{or} \quad \frac{3+\sqrt{3}}{\sqrt{3}-1} \times \frac{\sqrt{3}+1}{\sqrt{3}+1} = \frac{3\sqrt{3}+3+3+\sqrt{3}}{(\sqrt{3}-1)(\sqrt{3}+1)}$$

Allow  $2\sqrt{3}+3$ . It is for this expression and not for just  $p=3, q=2$  unless  $3+2\sqrt{3}$  is seen.

**Example of insufficient work:**

$$x\sqrt{3}-3 = x+\sqrt{3} \Rightarrow x\sqrt{3}-x = 3+\sqrt{3} \Rightarrow x(\sqrt{3}-1) = 3+\sqrt{3} \Rightarrow x = \frac{3+\sqrt{3}}{\sqrt{3}-1}$$

$$= \frac{3+\sqrt{3}}{\sqrt{3}-1} \times \frac{\sqrt{3}+1}{\sqrt{3}+1} = 3+2\sqrt{3}$$

**Scores M1M1A0**


**Note that:**

$$x\sqrt{3}-3 = x+\sqrt{3} \Rightarrow x\sqrt{3}-x = 3+\sqrt{3} \Rightarrow x(\sqrt{3}-1) = 3+\sqrt{3} \Rightarrow x = \frac{3+\sqrt{3}}{\sqrt{3}-1} = 3+2\sqrt{3}$$

**Scores M1M0A0**

**Alternatives for part (ii)**

<b>(ii)</b> <b>ALT 1</b>	$x\sqrt{3} - 3 = x + \sqrt{3} \Rightarrow (p + q\sqrt{3})\sqrt{3} - 3 = p + q\sqrt{3} + \sqrt{3} \Rightarrow p\sqrt{3} + 3q - 3 = p + \sqrt{3} + q\sqrt{3}$ $\Rightarrow p = q + 1, 3q - 3 = p$	M1
	$\Rightarrow p = q + 1, 3q - 3 = p \Rightarrow p = \dots, q = \dots$	M1
	$x = 3 + 2\sqrt{3}$	A1
<b>Notes:</b>		
<b>M1:</b> Substitutes $x = p + q\sqrt{3}$ , collects terms, compares coefficients and forms 2 equations in $p$ and $q$ It is for obtaining 2 equations of the form $\alpha p + \beta q = \gamma$ , $\alpha, \beta, \gamma \neq 0$		
<b>M1:</b> Solves simultaneously with evidence of algebra i.e. answers not just written down from a calculator. Not formally dependent but requires the solution of 2 equations of the form $\alpha p + \beta q = \gamma$ , $\alpha, \beta, \gamma \neq 0$		
<b>A1:</b> For $3 + 2\sqrt{3}$ . Allow $2\sqrt{3} + 3$ . It is for this expression and not for just $p = 3, q = 2$ unless $3 + 2\sqrt{3}$ is seen.		
<b>(ii)</b> <b>ALT 2</b>	$x\sqrt{3} - 3 = x + \sqrt{3} \Rightarrow 3x^2 - 6\sqrt{3}x + 9 = x^2 + 2\sqrt{3}x + 3$ $\Rightarrow 2x^2 - 8\sqrt{3}x + 6 = 0$	M1
	$x^2 - 4\sqrt{3}x + 3 = 0 \Rightarrow x = \frac{4\sqrt{3} \pm \sqrt{(4\sqrt{3})^2 - 4 \times 1 \times 3}}{2}$	M1
	$\Rightarrow x = \frac{4\sqrt{3} \pm \sqrt{36}}{2} \Rightarrow x = 2\sqrt{3} \pm 3$ $x = 2\sqrt{3} + 3$	A1
<b>Notes:</b>		
<b>M1:</b> Squares both sides. Must obtain a 3 term quadratic expression on both sides and collect terms to one side.		
<b>M1:</b> Solves a 3TQ of the form $px^2 + q\sqrt{3}x + r = 0$ by a correct non-calculator method.		
<b>A1:</b> For $3 + 2\sqrt{3}$ . Allow $2\sqrt{3} + 3$ . It is for this expression and not for just $p = 3, q = 2$ unless $3 + 2\sqrt{3}$ is seen. If any other answers are offered and clearly not rejected score A0.		
<b>(ii)</b> <b>ALT 3</b>	$x\sqrt{3} - 3 = x + \sqrt{3} \Rightarrow (x\sqrt{3} - 3)(x - \sqrt{3}) = (x + \sqrt{3})(x - \sqrt{3})$ $\Rightarrow \sqrt{3}x^2 - 6x + 3\sqrt{3} = x^2 - 3 \Rightarrow (\sqrt{3} - 1)x^2 - 6x + 3\sqrt{3} + 3 = 0$	M1
	$(\sqrt{3} - 1)x^2 - 6x + 3\sqrt{3} + 3 = 0 \Rightarrow x = \frac{6 \pm \sqrt{36 - 4(\sqrt{3} - 1)(3\sqrt{3} + 3)}}{2(\sqrt{3} - 1)}$ $= \frac{6 \pm \sqrt{36 - 24}}{2(\sqrt{3} - 1)} = \frac{6 \pm \sqrt{12}}{2(\sqrt{3} - 1)} = \frac{3 \pm \sqrt{3}}{\sqrt{3} - 1} = \frac{3 \pm \sqrt{3}}{\sqrt{3} - 1} \times \frac{\sqrt{3} + 1}{\sqrt{3} + 1}$	M1
	$= \frac{3\sqrt{3} + 3 \pm 3 \pm \sqrt{3}}{2} = 2\sqrt{3} + 3, \sqrt{3}$ $x = 2\sqrt{3} + 3$	A1
<b>Notes:</b>		
<b>M1:</b> Multiplies both sides by e.g. $(x - \sqrt{3})$ and collects terms to one side. Must obtain a 3 term quadratic expression on lhs and a 2 or 3 term quadratic expression on rhs.		
<b>M1:</b> Solves a 3TQ of the form $(A + B\sqrt{3})x^2 + Cx + D + E\sqrt{3} = 0$ , $A, B, C, D, E \neq 0$ by a correct non-calculator method <b>and</b> then attempts to rationalise the denominator as in the main scheme.		
<b>A1:</b> For $3 + 2\sqrt{3}$ . Allow $2\sqrt{3} + 3$ . It is for this expression and not for just $p = 3, q = 2$ unless $3 + 2\sqrt{3}$ is seen. If any other answers are offered and clearly not rejected score A0.		
		Total 6

Question	Scheme	Marks
3(a)		B1B1B1
		<b>(3)</b>

**Notes:**

**If there is any ambiguity regarding coordinates, the sketch takes precedence.**

**B1:** Same shape translated parallel to the  $x$ -axis with no evidence **from the  $x$ -intercepts** (if labelled) that the transformation is anything other than a translation. So the  $x$  intercepts (if labelled) must be 4 units apart. Ignore any coordinates on or near the maximum.


**B1:** Minimum at the origin. The origin does not have to be labelled.

**B1:** Passes through  $(-4, 0)$  or at least “reaches” this point from above or below.

Allow just  $-4$  marked in the correct place and condone  $(0, -4)$  as long as it is marked in the correct place.

Allow to score away from the sketch as long as their curve passes through (or reaches) this point but must be identified correctly as  $(-4, 0)$  and correspond to the sketch.


**See examples below:**



B1: Translation (the maximum at  $(-3, 2)$  can be ignored)

B1: Minimum at the origin


B1: Passes through  $(-4, 0)$



B0: The  $x$  intercepts are contradictory for a translation (not 4 units apart)

B0: No minimum at the origin


B1: Passes through  $(-4, 0)$



B1: Translation

B1: Minimum at the origin (the intention is clear)

B1: Passes through  $(-4, 0)$



B0: Incorrect shape

B1: Minimum at the origin

B1: Passes through  $(-4, 0)$

(b)		B1B1B1
		(3)
		<b>Total 6</b>

**Notes:**

**If there is any ambiguity regarding coordinates, the sketch takes precedence.**

**B1:** Reflection in the  $y$ -axis. This requires a minimum on the negative  $x$ -axis and a maximum to the left of the  $y$ -axis in quadrant 2. If the maximum is on the  $y$ -axis score B0.


The curve must continue upwards to the left of the  $y$ -axis to reach a maximum in quadrant 2.

The curve must extend into quadrant 4 as shown.

**B1:** Minimum at  $(-1, 0)$ . Allow just  $-1$  marked in the correct place and condone  $(0, -1)$  as long as it is marked in the correct place. Allow to score away from the sketch but must be identified correctly as  $(-1, 0)$  and correspond to the sketch.

**B1:** Passes through  $(0, 2)$  and passes through or touches  $(\frac{1}{3}, 0)$ . For both points, allow just the intercepts marked in the correct place and condone  $(2, 0)$  and/or  $(0, \frac{1}{3})$  as long as they are marked in the correct place. Allow to score away from the sketch but must be identified correctly as  $(0, 2)$  and  $(\frac{1}{3}, 0)$  and correspond to the sketch.


**See examples below:**



B0: Maximum in quadrant 2

B1: Minimum at  $(-1, 0)$


B1: Passes through  $(0, 2)$  and  $(\frac{1}{3}, 0)$



B0: Maximum on the  $y$ -axis (we would have tolerated the slight “pointy” maximum)

B0: Minimum at  $(-2, 0)$


B0: Does not pass through  $(0, 2)$  and  $(\frac{1}{3}, 0)$



B0: Not a reflection in the  $y$ -axis

B0: No minimum at  $(-1, 0)$

B1: Passes through  $(0, 2)$  and touches  $(\frac{1}{3}, 0)$



B0: Must be a curve – a reflection in the  $y$ -axis with the same shape.

B1: Minimum at  $(-1, 0)$

B1: Passes through  $(0, 2)$  and  $(\frac{1}{3}, 0)$

Question	Scheme	Marks
<b>4(a)</b>	$x^2 + kx - 9 = -3x^2 - 5x + k \Rightarrow 4x^2 + kx + 5x - 9 - k (= 0)$	M1
	$b^2 - 4ac = 0 \Rightarrow (k + 5)^2 - 4 \times 4(-9 - k) = 0$	M1
	$k^2 + 26k + 169 = 0^*$	A1*
		<b>(3)</b>
<b>(b)</b>	$k^2 + 26k + 169 = 0 \Rightarrow k = -13$	B1
	$k = -13 \Rightarrow 4x^2 - 8x + 4 = 0 \Rightarrow x = \dots$	M1
	$(1, -21)$	A1
		<b>(3)</b>
		<b>Total 6</b>

**Notes:**

**Mark (a) and (b) together**

(a)

**M1:** Equates both curves and collects terms to one side. Allow slips but there must be some attempt to collect terms to one side. The “= 0” may be implied by their attempt at the discriminant.

**M1:** Attempts to find the discriminant “ $b^2 - 4ac$ ” of a 3TQ. May be seen as an attempt at e.g.  $b^2 - 4ac = 0$  or equivalent e.g.  $b^2 = 4ac$  or may be seen embedded in an attempt at the quadratic formula.

It requires  $b$  and  $c$  both of the form  $pk + q$ ,  $p, q \neq 0$  and requires  $a$  as a constant.

If clearly a wrong formula/expression is used e.g. “ $b^2 + 4ac$ ” this scores M0

**A1\*:** Achieves the printed answer with no errors and sufficient working shown.

The “= 0” must appear at least once before the printed answer unless they start with  $b^2 = 4ac$ .

(b)

**B1:** Correct value of  $k$ . Condone  $x = -13$  if it is subsequently used as a value for  $k$ .

**M1:** Either:

- substitutes their value of  $k$  into their equation obtained by equating the curves and collecting terms from part (a) and solves a 3TQ for  $x$  by any correct method including a calculator **or**
- substitutes their  $k$  into the 2 given equations, equates, collects terms to one side and solves a 3TQ for  $x$  by any correct method including a calculator

**A1:** Correct coordinates. Allow as e.g.  $x = 1$ ,  $y = -21$

**Correct answer only in (b) scores 3/3**

Question	Scheme	Marks
<b>5(a)</b>	$\frac{1}{2} \times 6^2 \times 1.3 = \dots$	M1
	$= 23.4 \text{ (m}^2\text{)}$	A1
		(2)
<b>(b)</b>	$12.2^2 = 6^2 + 10.8^2 - 2 \times 6 \times 10.8 \cos(ABE)$	M1
	$\cos(ABE) = \frac{6^2 + 10.8^2 - 12.2^2}{2 \times 6 \times 10.8} \left( = \frac{19}{648} \right)$ $ABE = 1.54$	A1
		(2)
<b>(c)</b>	$\text{Area } ABE = \frac{1}{2} \times 10.8 \times 6 \sin(ABE)$	M1
	$\text{Area } BCD = \frac{1}{2} \times 6 \cos(\pi - 1.3 - "1.54") \times 6 \sin(\pi - 1.3 - "1.54")$ or e.g. $\text{Area } BCD = \frac{1}{2} \times 6 \sin(\pi - 1.3 - "1.54") \times \sqrt{6^2 - (6 \sin(\pi - 1.3 - "1.54"))^2}$	M1
	$\text{Total area} = 60.9 \text{m}^2$	A1
		(3)
		<b>Total 7</b>

**Notes:**

(a) **Marks for part (a) must be seen in part (a)**

**M1:** Attempts to evaluate  $\frac{1}{2} r^2 \theta$  with  $r = 6$  and  $\theta = 1.3$  or equivalent e.g.  $\frac{\theta}{360} \times \pi \times 6^2$  with  $\theta = 1.3 \times \frac{180}{\pi}$  which may be implied by use of e.g.  $74^\circ$  or  $75^\circ$

**A1:** Correct value. Allow equivalent values e.g.  $\frac{117}{5}$ ,  $23\frac{2}{5}$ .

Condone lack of units but if any are given they should be correct. Correct answer only scores both marks.

(b)

**M1:** Correct numerical application of the cosine rule for the required angle.

**A1:** For awrt 1.54

(c)

**Allow equivalent work in degrees.**

**M1:** Correct method for area  $ABE$  e.g.  $\frac{1}{2} \times 10.8 \times 6 \sin(\text{their } ABE)$

**M1:** Correct method for area  $BCD$

Must be a correct method including the  $\frac{1}{2}$

Note that  $CD$  may be found using the sine rule e.g.  $\frac{6}{\sin \frac{\pi}{2}} = \frac{CD}{\sin(\pi - 1.3 - "1.54")} \Rightarrow CD = \dots$

Note that  $BC$  may be found using the sine rule e.g.  $\frac{6}{\sin \frac{\pi}{2}} = \frac{BC}{\sin\left(1.3 + "1.54" - \frac{\pi}{2}\right)} \Rightarrow BC = \dots$

Having found  $CD$  or  $BC$  as above,  $BC$  or  $CD$  may be found using Pythagoras.

In such cases the trigonometry must be correct for their values.

Do **not** allow mixing of degrees and radians e.g.  $180 - 1.3 - "1.54"$  for  $\pi - 1.3 - "1.54"$

**A1:** Correct total area. Allow awrt 60.9 (Condone lack of units)

**Some values for reference:**

Angle  $DBC = 0.300\dots(17.1\dots^\circ)$ , Angle  $BDC = 1.27\dots(72.7\dots^\circ)$ ,  
Area  $ABE = 32.38\dots$ , Area  $BCD = 5.083\dots$ ,  $CD = 1.77\dots$ ,  $BC = 5.73\dots$

Question	Scheme	Marks
<b>6(a)</b>	$y-5x=75, y=2x^2+x-21$ $\Rightarrow 2x^2+x-21=5x+75$ $\Rightarrow 2x^2-4x-96=0 \text{ or e.g. } x^2-2x-48=0$ <p style="text-align: center;">or</p> $\Rightarrow y=2\left(\frac{y-75}{5}\right)^2+\frac{y-75}{5}-21$ $\Rightarrow 2y^2-320y+10350=0 \text{ or e.g. } y^2-160y+5175=0$	M1
	$x^2-2x-48=0 \Rightarrow (x-8)(x+6)=0 \Rightarrow x=-6, 8$	dM1
	$x=-6 \Rightarrow y=45 \text{ or } x=8 \Rightarrow y=115$	dM1
	$P(-6, 45) \text{ and } Q(8, 115)$	A1
		<b>(4)</b>
<b>(b)</b>	e.g. $y \leq 2x^2+x-21, y-5x \leq 75, y \geq 0, x \leq -3.5$  $x \leq a \text{ where } -3.5 \leq a < 3$ (or $a \leq x \leq b \text{ where } a \leq -15, -3.5 \leq b \leq 3$ )	M1A1A1
		<b>(3)</b>
		<b>Total 7</b>

**Notes:**

(a) **Note that solutions relying on calculator technology are not acceptable so a complete method must be shown.**

**M1:** Equates the line and the curve and rearranges to obtain a 3TQ in  $x$

Alternatively eliminates  $x$  to obtain a 3TQ in  $y$

**dM1:** Solves their 3TQ to obtain at least one value for  $x$  or  $y$  by an acceptable method e.g. factorisation, formula or completing the square. They cannot just state the roots without seeing a correct line of intermediate working.

We will condone e.g.  $2x^2-4x-96=0 \Rightarrow x = \frac{4 \pm \sqrt{(-4)^2 - 4(2)(-96)}}{2 \times 2} = -6, 8$

But e.g.  $2x^2-4x-96=0 \Rightarrow x = \frac{4 \pm \sqrt{-4^2 - 4(2)(-96)}}{2 \times 2} = -6, 8$  scores M0 as it suggests an incorrect

formula but allow recovery e.g.  $2x^2-4x-96=0 \Rightarrow x = \frac{4 \pm \sqrt{-4^2 - 4(2)(-96)}}{2 \times 2} = \frac{4 \pm \sqrt{784}}{4} = -6, 8$  and

condone a missing trailing bracket e.g.  $x = \frac{4 \pm \sqrt{(-4)^2 - 4(2)(-96)}}{2 \times 2} = -6, 8$

If they factorise do **not** accept  $2x^2-4x-96=0 \Rightarrow (x+6)(x-8)=0 \Rightarrow x=-6, 8$

**Depends on the previous method mark.**

**dM1:** Uses at least one  $x$  value to find at least one  $y$  value (or vice versa) using either of the given equations.

**Depends on the FIRST method mark and depends on having solved a 3TQ in  $x$  or  $y$ .**

May be implied by their values. You may need to check (for both equations).

**A1:** All correct. Must be correctly paired but ignore any labelling regarding  $P$  and  $Q$ , just look for the correct paired values. This may be explicit as a coordinate pair or implied by e.g.  $x=8, y=115$ .

Condone coordinates written without brackets e.g.  $-6, 45$

**Depends on all previous marks so 1011 is not possible.**

**Candidates who show no working for solving their 3TQ can score a maximum 1010**

**Correct answers with no working scores no marks.**

**(b) Allow strict or non-strict inequalities and allow a mixture for the first 2 marks.**

**M1:** Obtains at least 2 of the required inequalities e.g. 2 of:

$$y \leq 2x^2 + x - 21, y - 5x \leq 75, y \geq 0, x \leq a \text{ where } -3.5 \leq a < 3$$

Allow equivalents e.g.  $y < 5x + 75$

Allow e.g.  $0 \leq y \leq 2x^2 + x - 21$  or e.g.  $0 < y < 5x + 75$  each of which would count as 2 correct inequalities.

Do **not** allow e.g.  $R \leq 2x^2 + x - 21$

**A1:** Any **3** of the 4 required inequalities.

**A1:** All correct and consistent. For consistency, the inequalities for  $y$  must be either all strict or all non-strict.

The restriction on  $x$  will depend on their chosen  $x$  value. For example

$$y \geq 0, y \leq 5x + 75, y \leq 2x^2 + x - 21, x < 0 \text{ scores full marks}$$

$$y \geq 0, y \leq 5x + 75, y \leq 2x^2 + x - 21, x < -3.5 \text{ scores M1A1A0}$$

If there are extra **incorrect** inequalities then score A0



Question	Scheme	Marks
<b>7(a)(i)</b>	$f(x) = 2x^3 - kx^2 + 14x + 24 \Rightarrow (f'(x) =) 6x^2 - 2kx + 14$	M1A1
	$(f''(x) =) 12x - 2k$	A1ft
<b>(ii)</b>		<b>(3)</b>
	$6x^2 - 2kx + 14 = 12x - 2k \Rightarrow 6(5)^2 - 2k(5) + 14 = 12(5) - 2k \Rightarrow k = \dots$	M1
	$k = 13$	A1
<b>(b)</b>		<b>(2)</b>
	$k = 13 \Rightarrow 6x^2 - 38x + 40 = 0 \Rightarrow x = \dots$	M1
	$x = \frac{4}{3} \Rightarrow y = 12\left(\frac{4}{3}\right) - 2 \times 13$ or $y = 6\left(\frac{4}{3}\right)^2 - 2 \times 13 \times \left(\frac{4}{3}\right) + 14$	M1
	$x = \frac{4}{3}, y = -10$	A1
		<b>(3)</b>
		<b>Total 8</b>

**Notes:**

**In part (a), penalise the inclusion of “+ c” on either derivative only once on the first occurrence.**

(a)(i)

**M1:** For  $x^3 \rightarrow \dots x^2$  or  $x^2 \rightarrow \dots x$  or  $14x \rightarrow 14$

**A1:** Correct simplified first derivative. The “ $f'(x) =$ ” is not required.

Do not allow  $x^1$  for  $x$ .

Isw can be applied once a correct simplified derivative is seen.

(ii)

**A1ft:** Correct simplified second derivative. Follow through their first derivative

The “ $f''(x) =$ ” is not required.

Isw can be applied once a correct simplified second derivative is seen. **Must follow M1**

(b)

**M1:** Sets  $f'(x) = f''(x)$ , substitutes  $x = 5$  and solves a linear equation in  $k$  to find a value for  $k$ .

May substitute  $x = 5$  into  $f'(x)$  and  $f''(x)$  first, then equate and solve for  $k$ .

**A1:**  $k = 13$

(c)

**M1:** Substitutes their  $k$  into the equated  $f'(x) = f''(x)$  and solves the resulting 3TQ to obtain at least one value for  $x$  by any means including a calculator.

**M1:** Substitutes their  $x$  (not 5) and their  $k$  into  $f'(x)$  or  $f''(x)$  to find a value for  $y$ .

May be implied by their values so you may need to check.

**A1:** Correct coordinates. May be seen as  $x = \dots, y = \dots$  or as a coordinate pair but condone missing brackets

e.g.  $\frac{4}{3}, -10$ . Award this mark as soon as the correct values are seen.

Allow a “made up”  $k$  in (c).

Question	Scheme	Marks
8(a)		B1B1B1
		(3)
(b)	$x(4 - x^2) = \frac{A}{x} \Rightarrow 4x^2 - x^4 = A$ $\Rightarrow x^4 - 4x^2 + A = 0^*$	B1*
		(1)
(c)	$A > 0$	B1
	$b^2 = 4ac \Rightarrow 16 = 4A \Rightarrow A = \dots$	M1
	$0 < A < 4$	A1
		(3)
		<b>Total 7</b>

**Notes:**

(a)

**B1:** Negative cubic curve in any position. Must have one maximum and one minimum.

**B1:** Positive or negative cubic curve **passing through** (not touching) the origin.

The origin does not have to be labelled.

**B1:** Cuts the  $x$ -axis at  $x = 2$  and  $x = -2$ . May be as shown or as  $(2, 0)$  and  $(-2, 0)$ .


Must cut so extends beyond the  $x$ -axis from above or below but **not** a turning point.

Condone  $(0, 2)$  and  $(0, -2)$  as long as they are in the correct positions.


Allow away from the sketch but must be  $(2, 0)$  and  $(-2, 0)$ .

If there is any ambiguity, the sketch takes precedence.

**Examples:**



100




110




010




010




001




110



011



010



000

(b)

**B1\*:** Equates the curves and shows sufficient working to obtain the printed answer with no errors.

Note that  $x(4-x^2)$  may be expanded first before equating which is fine.

There must be at least one intermediate step between  $x(4-x^2) = \frac{A}{x}$  and the printed answer.

As a minimum accept  $x(4-x^2) = \frac{A}{x} \Rightarrow 4x - x^3 = \frac{A}{x} \Rightarrow x^4 - 4x^2 + A = 0$

(c)

**B1:** For  $A > 0$  as one boundary. May be seen embedded in their final answer.  $x > 0$  is B0.

**M1:** Attempts  $b^2 = 4ac$  or equivalent e.g.  $b^2 - 4ac = 0$  with  $a = 1$ ,  $b = -4$  and  $c = A$  to obtain a value for  $A$ .

Condone use of an inequality rather than “=” to obtain a value or inequality for  $A$  e.g.  $b^2 - 4ac > 0$  or

$b^2 - 4ac < 0$  leading to  $A > \dots$  or  $A < \dots$

May be implied by e.g.  $A = 4$ ,  $A < 4$ ,  $A > 4$  etc.

**A1:** Correct range for  $A$ . Allow equivalents e.g. “ $A > 0$  and  $A < 4$ ”, “(0, 4)”

but not “ $A > 0$  or  $A < 4$ ”, “ $A > 0, A < 4$ ”

**Correct answer with no working scores 3/3**

**Partially correct answer with no working e.g.  $0 < A \leq 4$  can score B1 M1 (implied) A0**

Question	Scheme	Marks	
<b>9(a)</b>	$m = \frac{7-2}{15-4} \left( = \frac{5}{11} \right)$	M1	
	$y-2 = \frac{5}{11}(x-4) \quad \text{or} \quad y-7 = \frac{5}{11}(x-15)$ or $y = \frac{5}{11}x + c \Rightarrow 2 = \frac{5}{11} \times 4 + c \Rightarrow c = \dots \left( \frac{2}{11} \right)$	M1	
	$5x - 11y + 2 = 0$	A1	
		<b>(3)</b>	
<b>(a) ALT</b>	$y = mx + c \Rightarrow \begin{cases} 2 = 4m + c \\ 7 = 15m + c \end{cases}$	M1	
	$\begin{cases} 2 = 4m + c \\ 7 = 15m + c \end{cases} \Rightarrow m = \dots \left( \frac{5}{11} \right), \quad c = \dots \left( \frac{2}{11} \right)$	M1	
	$5x - 11y + 2 = 0$	A1	
<b>(b)</b>	$(15-x)^2 + (7-2)^2 = (5\sqrt{5})^2$ or e.g. $\sqrt{(15-x)^2 + (7-2)^2} = 5\sqrt{5}$ or e.g. $(5\sqrt{5})^2 = 5^2 + CD^2 \text{ oe}$		M1
	Way 1: $(15-x)^2 + 25 = 125 \Rightarrow (15-x)^2 = 100 \Rightarrow x = \dots$ or e.g.		M1
	Way 2: $125 = 25 + CD^2 \Rightarrow CD^2 = 100 \Rightarrow CD = 10 \Rightarrow x = \dots$		
	$(5, 2) \text{ or } (25, 2)$		A1
	$(5, 2) \text{ and } (25, 2)$		A1
		<b>(4)</b>	
<b>(c)</b>	$\text{Area} = \frac{1}{2} ("5" - 4) \times 5$	M1	
	$= \frac{5}{2}$	A1	
		<b>(2)</b>	
		<b>Total 9</b>	

**Notes:**

(a)

**M1:** Correct method for the gradient

**M1:** For a correct straight line method using their gradient and point A or point B with x and y correctly placed.

Alternatively uses  $y = mx + c$  and reaches  $c = \dots$

Note that  $\frac{y-2}{7-2} = \frac{x-4}{15-4}$  scores both M marks.


**A1:** Correct equation in the required form including the “= 0”. Allow any integer multiple of this equation.

**ALT**

**M1:** Substitutes  $x = 4, y = 2$  and  $x = 15, y = 7$  into the general equation  $y = mx + c$  to obtain 2 equations in  $m$  and  $c$ .

**M1:** Solves 2 equations of the form  $\alpha = m\beta + c$  to find values for  $m$  and  $c$ .

**A1:** Correct equation in the required form including the “= 0”. Allow any integer multiple of this equation.



(b)

**M1:** Correct application of Pythagoras for triangle  $BCD$  (see diagram above)

The statements shown are sufficient for this mark but condone missing brackets e.g.  $5\sqrt{5}^2$  for  $(5\sqrt{5})^2$  as long as the intention is clear.

**M1:** A complete, correct and full method to obtain at least **one**  $x$  coordinate for  $C$ .

For **Way 1** this mark is for processing an equation of the form:  $(15-x)^2 + \alpha^2 = (5\sqrt{5})^2$  where  $0 < \alpha^2 < 125$  to obtain at least one value for  $x$ .

For **Way 2** this mark is for processing an equation of the form:  $(5\sqrt{5})^2 = CD^2 + \alpha^2$  where  $0 < \alpha^2 < 125$  to obtain at least one value for  $CD$  and then attempts at least one value of  $x$  either  $15 - CD$  or  $15 + CD$ . Condone the use of  $x$  (or any other letter) for  $CD$ .

The work for this mark must be correct so e.g.  $\sqrt{(15-x)^2 + (7-2)^2} = 5\sqrt{5} \Rightarrow (15-x) + (7-2) = 5\sqrt{5}$  or e.g.  $(5\sqrt{5})^2 = CD^2 + \alpha^2 \Rightarrow 5\sqrt{5} = CD + \alpha$  scores M0. If a 3TQ is being solved, the usual rules apply.

**A1:** At least one correct position for  $C$ .

May be seen as a coordinate pair with or without brackets or as e.g.  $x = 5, y = 2$

**A1:** Both correct pairs of coordinates.

May be seen as coordinate pairs with or without brackets or as e.g.  $x = 5, y = 2$

**Note that some candidates think the equation for  $l_2$  is  $y = 4$  rather than  $y = 2$**

**This scores a maximum of 1 mark in part (b) and will look like this:**

$$(15-x)^2 + (7-4)^2 = (5\sqrt{5})^2 \Rightarrow (15-x)^2 + (7-4)^2 = 125 - 9 = 116$$

$$(15-x) = \pm 2\sqrt{29} \Rightarrow x = 15 \pm 2\sqrt{29}$$

or

$$(5\sqrt{5})^2 = 3^2 + CD^2 \Rightarrow CD^2 = 116 \Rightarrow CD = 2\sqrt{29}$$

$$\Rightarrow x = 15 \pm 2\sqrt{29}$$

This scores M0M1A0A0

(c)

**M1:** Applies a correct triangle area for one of their points of the form  $(k, 2)$ ,  $k \neq 15$ . May be implied.

This mark is for e.g.  $\frac{1}{2}(k-4) \times 5$  where  $k$  is one of their calculated  $x$  coordinates from part (b)

or for  $\frac{1}{2}(11 \pm CD) \times 5$  where  $CD$  is their calculated value from part (b)

May see shoelace method:  $\frac{1}{2} \begin{vmatrix} 4 & 15 & "5" & 4 \\ 2 & 7 & 2 & 2 \end{vmatrix} = \frac{1}{2} |4 \times 7 + 15 \times 2 + "5" \times 2 - 2 \times 4 - 7 \times "5" - 2 \times 15|$

Allow other correct methods for the area e.g. using trigonometry.

Their  $C$  must be of the form  $(k, 2)$  and a correct formula used or implied.

**A1:** Correct minimum area. Must be 2.5 **not** awrt 2.5 (e.g. if they use trigonometry and get an inexact answer)

Question	Scheme	Marks
<b>10(a)</b>	$f'(4) = 6(4) - \frac{7 \times 14}{4} = -\frac{1}{2}$	B1
	$m_T = -\frac{1}{2} \Rightarrow m_N = \frac{-1}{-\frac{1}{2}}$	M1
	$y - 12 = 2(x - 4)$ or $y = mx + c \Rightarrow y = 2x + c \Rightarrow 12 = 2(4) + c \Rightarrow c = \dots$	M1 (A1 on ePen)
	$y = 2x + 4$	A1
		(4)
<b>(b)</b>	$\dots \frac{(2x-1)(3x+2)}{2\sqrt{x}} = \dots \frac{6x^2+x-2}{2\sqrt{x}} = \dots 3x^{\frac{3}{2}} + \frac{1}{2}x^{\frac{1}{2}} - x^{-\frac{1}{2}}$	M1
	$f(x) = \frac{6x^2}{2} - \frac{6}{5}x^{\frac{5}{2}} - \frac{1}{3}x^{\frac{3}{2}} + 2x^{\frac{1}{2}} (+c)$ or e.g. $f(x) = \frac{6x^2}{2} - \frac{3}{\frac{5}{2}}x^{\frac{5}{2}} - \frac{1}{\frac{3}{2}}x^{\frac{3}{2}} + \frac{x^{\frac{1}{2}}}{\frac{1}{2}} (+c)$ or e.g. $f(x) = \frac{6x^2}{2} - \left( \frac{3}{\frac{5}{2}}x^{\frac{5}{2}} + \frac{1}{\frac{3}{2}}x^{\frac{3}{2}} - \frac{x^{\frac{1}{2}}}{\frac{1}{2}} \right) (+c)$	M1A1A1
	$12 = \frac{6(4)^2}{2} - \frac{6}{5}(4)^{\frac{5}{2}} - \frac{1}{3}(4)^{\frac{3}{2}} + 2(4)^{\frac{1}{2}} + c \Rightarrow c = \dots$	M1
	$(f(x) =) 3x^2 - \frac{6}{5}x^{\frac{5}{2}} - \frac{1}{3}x^{\frac{3}{2}} + 2x^{\frac{1}{2}} + \frac{16}{15}$	A1
		(6)
		<b>Total 10</b>

**Notes:**

(a)

**B1:** Correct gradient at  $P$  (may be implied)

**M1:** Attempts to use the perpendicular gradient rule to find the normal gradient.

Look for e.g.  $m_N = \frac{-1}{m_T}$  or e.g.  $m_T \times m_N = -1 \Rightarrow -\frac{1}{2}m_N = -1 \Rightarrow m_N = \dots$

May be implied by their normal gradient.

**M1(A1 on Epen):** Attempts the equation of the normal using a “changed” gradient with  $x = 4$  and  $y = 12$  correctly placed.

Alternatively uses  $y = mx + c$  with a “changed” gradient with  $x = 4$  and  $y = 12$  to find a value for  $c$ .

**A1:** Correct equation in the required form

(b) **Allow work for part (b) seen in part (a) to score in (b) provided it is seen or used in (b)**

**M1:** Expands the numerator of the fraction and attempts to split.

Score for one correct index achieved from correct work e.g.  $\frac{\dots x^2}{\sqrt{x}} \rightarrow \dots x^{\frac{3}{2}}$  or  $\frac{\dots x}{\sqrt{x}} \rightarrow \dots x^{\frac{1}{2}}$  or  $\frac{\dots}{\sqrt{x}} \rightarrow \dots x^{-\frac{1}{2}}$

**M1:** Attempts to integrate a fractional power. E.g.  $\dots x^{\frac{3}{2}} \rightarrow \dots x^{\frac{5}{2}}$  or  $x^{\frac{1}{2}} \rightarrow \dots x^{\frac{3}{2}}$  or  $x^{-\frac{1}{2}} \rightarrow \dots x^{\frac{1}{2}}$  etc.

Do not allow this mark for an attempt to integrate the  $\sqrt{x}$  in the denominator.

**A1:** Any 2 correct terms simplified or unsimplified.

**A1:** All correct simplified or unsimplified. The “+ c” is not required here.

**M1:** Uses  $x = 4$  and  $y = 12$  following an attempt to increase at least one power  $x$  by 1 to find a value for their  $c$ .

Their  $c$  may not be fully evaluated but must be a numerical expression.

**A1:** Correct simplified form. The “f(x) = ” is not required just look for the correct expression.

Accept equivalent simplified forms and allow  $(f(x) =) 3x^2 - \left( \frac{6}{5}x^{\frac{5}{2}} + \frac{1}{3}x^{\frac{3}{2}} - 2x^{\frac{1}{2}} \right) + \frac{16}{15}$

Apply isw if applicable unless they multiply all terms by e.g. 15

Question	Scheme	Marks
<b>11(a)</b>	$x = \frac{5\pi}{2}$ or $y = 12$	B1
	$x = \frac{5\pi}{2}$ and $y = 12$	B1
		(2)
<b>(b)</b>	$x = \frac{3\pi}{2}$ or $y = -21$	B1
	$x = \frac{3\pi}{2}$ and $y = -21$	B1
		(2)
<b>(c)(i)</b>	$(A =) -12$	B1
<b>(ii)</b>	$(B =) \frac{5\pi}{4}$	B1
		(2)
		<b>Total 6</b>

**Notes:**

**Look for answers written in the body of the question or on the sketch.**

**In all cases, answers written in the body of the script take precedence over answers written on the diagram.**

**Angles and numbers must be written as single terms e.g. not  $2\pi + \frac{\pi}{2}$  for  $\frac{5\pi}{2}$**

**Penalise the use of degrees once the first time it occurs**

e.g. (a)  $(450^\circ, 12)$  (b)  $(270^\circ, -21)$  (c)  $A = -12, B = 225^\circ$  scores (a) B1B0 (b) B1B1 (c) B1B1

**The degrees symbol is not required**

**If both coordinates in (a) and (b) are correct but the wrong way round score as B1B0 each time**

e.g. (a)  $\left(12, \frac{5\pi}{2}\right)$  (b)  $\left(-21, \frac{3\pi}{2}\right)$  scores (a) B1B0 (b) B1B0

**If the coordinates in (a) or (b) are the wrong way round and only one is correct score as B0B0 each time**

**If the coordinates are the wrong way round and correct in degrees in (a) and (b)**

e.g. (a)  $(12, 450^\circ)$  (b)  $(-21, 270^\circ)$  award (a) B0B0 (b) B1B0

**The degrees symbol is not required**



(a)

**B1:** One coordinate correct.

May be seen as  $x = \dots$  or  $y \dots$  or embedded in a coordinate pair e.g.  $\left(\frac{5\pi}{2}, 12\right)$  or in a vector  $\begin{pmatrix} \frac{5\pi}{2} \\ 12 \end{pmatrix}$

**B1:** Both coordinates correct.

May be seen as  $x = \dots$  and  $y \dots$  or as a coordinate pair e.g.  $\left(\frac{5\pi}{2}, 12\right)$  or as a vector  $\begin{pmatrix} \frac{5\pi}{2} \\ 12 \end{pmatrix}$

(b)

**B1:** One coordinate correct.

May be seen as  $x = \dots$  or  $y \dots$  or embedded in a coordinate pair e.g.  $\left(\frac{3\pi}{2}, -21\right)$  or in a vector  $\begin{pmatrix} \frac{3\pi}{2} \\ -21 \end{pmatrix}$

**B1:** Both coordinates correct.

May be seen as  $x = \dots$  and  $y \dots$  or as a coordinate pair e.g.  $\left(\frac{3\pi}{2}, -21\right)$  or as a vector  $\begin{pmatrix} \frac{3\pi}{2} \\ -21 \end{pmatrix}$

(c)(i)

**B1:** Correct value for  $A$ . The “ $A =$ ” is not required but it must be clear it is the answer to (i) or is the value of  $A$ .

(ii)

**B1:** Correct value for  $B$ . The “ $B =$ ” is not required but it must be clear it is the answer to (ii) or is the value of  $B$ .

If candidates write the values as a coordinate pair e.g.  $\left(\frac{5\pi}{4}, -12\right)$  or  $\left(-12, \frac{5\pi}{4}\right)$  so it is not clear which value is which, score B0B0

